

## Bilateral avascular necrosis of both hips and both shoulders in a case with systemic lupus...

Last Updated Wednesday, 28 March 2007

APLAR Journal of Rheumatology

Emad, Y., et al. - A 38-year-old female patient was diagnosed as a case of systemic lupus erythematosus (SLE) in 1994. Her initial presentation was nephritis which remitted on combination of steroid, azathioprine and pulse cyclophosphamide therapy. One year later the patient developed bilateral avascular necrosis (AVN) both hips and underwent bilateral hip replacement...It is suggested that in SLE patients with positive lupus anticoagulant and negative antiphospholipid antibodies, testing for reactivity against beta2 glycoprotein 1 is mandatory

APLAR Journal of Rheumatology

Emad, Y., et al. - A 38-year-old female patient was diagnosed as a case of systemic lupus erythematosus (SLE) in 1994. Her initial presentation was nephritis which remitted on combination of steroid, azathioprine and pulse cyclophosphamide therapy. One year later the patient developed bilateral avascular necrosis (AVN) both hips and underwent bilateral hip replacement...It is suggested that in SLE patients with positive lupus anticoagulant and negative antiphospholipid antibodies, testing for reactivity against beta2 glycoprotein 1 is mandatory

Read article: <http://www.mdlinx.com/OrthoLinx/xml-article.cfm/1793754>.