

## Role of Maximal Heart Rate and Arterial O<sub>2</sub> Saturation on the Decrement of V̇O<sub>2</sub>max in Moderate..

Last Updated Wednesday, 28 March 2007

International Journal of Sports Medicine

P. Mollard, et al. - We aimed to evaluate 1) the altitude where maximal heart rate (HR<sub>max</sub>) decreases significantly in both trained and untrained subjects in moderate acute hypoxia, and 2) if the HR<sub>max</sub> decrease could partly explain the drop of V̇O<sub>2</sub>max...Thus, in moderate acute hypoxia, the reduction in SaO<sub>2</sub> is the primary factor to explain the drop of V̇O<sub>2</sub>max in trained and untrained subjects

International Journal of Sports Medicine

P. Mollard, et al. - We aimed to evaluate 1) the altitude where maximal heart rate (HR<sub>max</sub>) decreases significantly in both trained and untrained subjects in moderate acute hypoxia, and 2) if the HR<sub>max</sub> decrease could partly explain the drop of V̇O<sub>2</sub>max...Thus, in moderate acute hypoxia, the reduction in SaO<sub>2</sub> is the primary factor to explain the drop of V̇O<sub>2</sub>max in trained and untrained subjects

Read article: <http://www.mdlinx.com/OrthoLinx/xml-article.cfm/1788841>.