

A Complete Expression Profile of Matrix-Degrading Metalloproteinases in Dupuytren's Disease

Last Updated Wednesday, 28 March 2007

Journal of Hand Surgery

Phillip Johnston MB BChir, et al. - Conclusions: This study measured the expression of all MMP, ADAMTS, and TIMP genes in DD. Contraction and fibrosis may result from: (1) increased collagen biosynthesis mediated by increased ADAMTS-14; (2) an increased level of TIMP-1 blocking MMP-1 and MMP-13 mediated collagenolysis; and (3) contraction enabled by MMP-14 mediated pericellular collagenolysis (and potentially MMP-7), which may escape inhibition by TIMP-1

Journal of Hand Surgery

Phillip Johnston MB BChir, et al. - Conclusions: This study measured the expression of all MMP, ADAMTS, and TIMP genes in DD. Contraction and fibrosis may result from: (1) increased collagen biosynthesis mediated by increased ADAMTS-14; (2) an increased level of TIMP-1 blocking MMP-1 and MMP-13 mediated collagenolysis; and (3) contraction enabled by MMP-14 mediated pericellular collagenolysis (and potentially MMP-7), which may escape inhibition by TIMP-1

Read article: <http://www.mdlinx.com/OrthoLinx/xml-article.cfm/1793681>.